

RECOMMENDED PRESERVATION & HOLDING TIMES (RHT) FOR WATERS

www.envirolab.com.au

ADELAIDE:	(08) 7087 6800	BRISBANE: (07) 3266 9532	DARWIN: 0477 012 027
MELBOURNE	(03) 9763 2500	SYDNEY: (02) 9910 6200	PERTH: (08) 9317 2505

Test	Bottle	Volume	Preservation RHT (Bottles are colour coded, see labels)		Reference
INORGANICS,	METALS AI	ND PHYSI			
Alkalinity / Acidity	P/G	100 ml	Cool to <u><</u> 6°C	14 d	АРНА
AOX	G	500 ml	pH <2 (HNO₃) + Cool to <u><</u> 6°C + Dark	3 d	AS
BOD	P/G	500 ml	Cool to <u><</u> 6°C	2 d / 1 month	APHA / ISO
				frozen	
Bromide	P/G	50 ml	Cool to <u><</u> 6°C	28 d	ISO
Bromate	Р	100 ml	Cool to <u><</u> 6°C 50mg/L EDA	28 d	ISO
Carbon - TOC/DOC	P/G	50 ml	pH <2 (H₂SO₄ or HCI) + Cool to <u><</u> 6°C	28 d	APHA
Carbon - TOC/DOC	P/G	50 ml	Cool to <u>≤</u> 6°C + Dark	28 d	USEPA
Carbon Dioxide (free	Р	500ml	Cool to <u><</u> 6°C (field measurement preferred)	15 min	APHA
+ total)					
Chlorine-Residual	P/G	20 ml	Analyse immediately	15 min	АРНА
Chlorite ^{\$} /Chlorate	Р	100 ml	Cool to <u><</u> 6°C 50mg/L EDA	14/28 d	USEPA
Chloride	P/G	100 ml	Nil	28 d	APHA
COD	P/G	100 ml	pH <2 (H₂SO₄) + Cool to <u><</u> 6°C	28 d / 6m	APHA / ISO
Colour	P/G	100 ml	Cool to <u><</u> 6°C + Dark	2 d / 5 d	APHA / ISO
Conductivity	P/G	100 ml	Cool to <u><</u> 6°C	28 d	USEPA
Cyanides/Cyanates	P/G	100 ml	pH (>12 NaOH) + Cool to <u><</u> 6°C + Dark	14 d	АРНА
Dust Deposition	G (Winchester)	4L	Copper Sulphate (10mL)	30 d	AS
Ferrous Iron	P/G	100 ml	Filter, pH <2 (HCI), no headspace	1 d / 7 d	AS / ISO
Fluoride	P (not PTFE)	50 ml	Nil	28 d	AS
Formaldehyde	Vial	2x40mL	Cool to <u><</u> 6°C	7 d / 14 d	ISO / USEPA
Hexavalent Cr	P/G	100 ml	Filter, cool to <u><</u> 6°C	1 d	AS
Hexavalent Cr	P/G	100 ml	Filter, pH 8-9.5 (NaOH) + Cool to <u><</u> 6°C	28 d	USEPA m1636 / APHA
lodide/lodine	P/G	50 ml	Cool to <u><</u> 6°C	28 d	APHA / in house
[all forms of lodine]					
MBAS - Surfactants	G	100 ml	Cool to <u><</u> 6°C	2 d	APHA
Metals (inc. Cations)	Р	50 ml	pH <2 (HNO ₃)	6 m	USEPA / APHA
Mercury	Р	50 ml	pH <2 (HNO₃)	28 d	USEPA/ APHA
Metals (inc. Cations)			Nil – Orange label bottle for airfreight will be lab acidified in <14days	6 m	USEPA / APHA
Mercury	Р	50ml	Nil – Orange label bottle for airfreight will be lab acidified in <14days	28 d	USEPA/ APHA
Anions/Cations/	P/G	500 ml	Anions (F ⁻ , Cl ⁻ , SO4 ²⁻ , Alkalinity), Cations (Ca, K, Mg, Na),	See individual	See individual tests
Nutrients/			Nutrients (NOx, NH ₃ , o-PO ₄), EC/pH, TDS and TSS can be	tests	
Physical Tests			analysed from one 500mL unpreserved sample.		
N – Ammonia	P/G	20 ml	pH <2 (H₂SO₄) + Cool to <u><</u> 6°C	28 d	APHA
N – Ammonia	P/G	20 ml	Site filter & cool / Site filter & freeze	1d / 28 d	AS / ISO
N – Nitrate + Nitrite	P/G	20 ml	Cool to <u><</u> 6°C / Filter on site & cool to <u><</u> 6°C	2 d / 4 d	APHA / ISO
N – Nitrate	P/G	20 ml	Filter on site & freeze	28 d	AS / ISO
N – Total and N-TKN	P/G	100 ml	pH <2 (H₂SO₄) + Cool to <u><</u> 6°C	28 d	APHA
N – Total and N-TKN	P/G	100 ml	Freeze	28 d	ISO
Oil & Grease	G	500 ml	pH <2 (H₂SO₄ or HCI) + Cool to <u><</u> 6°C	28 d	АРНА
Perchlorate	Р	100 mL	Cool to <u><</u> 6°C	28 d	USEPA
рН	P/G	100 ml	 Cool to <u><</u> 6°C (field measurement preferred)	15min / 6 hrs / 1d	APHA / AS / ISO
Perchlorate	Р	100 ml	Cool to <u><</u> 6°C 50mg/L EDA (or unpreserved) [®]	14/28 d	USEPA
Phenolics - total	P/G	100 ml	pH <2 (H₂SO₄) + Cool to ≤6°C	28 d	АРНА
Phosphorus - total	P/G	50 ml	pH <2 (HNO₃)	28 d	АРНА
Phosphate as P	P/G	20 ml	Filter & Cool to <6°C / Filter & Freeze	2d / 28 d	APHA / AS / ISO



	Bottle	Volume	Preservation	RHT	Reference
INORGANICS.	METALS A	ND PHYS	(Bottles are colour coded, see labels) ICAL TESTS (continued)		
Reactive Silica	P/G	250ml	Cool to <6°C & Filter (preferably on site)	28 d	АРНА
Settleable Solids	P	1000ml	none	2 d	АРНА
Speciated Arsenic	Р	125 ml	Field filter, pH <2 (few drops of HCI) + Cool to <6°C	ASAP, 28 d max	USEPA
Specific Gravity	P	50 ml	Cool to <6°C	2 d	AS / APHA
Sugar	P/G	100 ml	Cool to <6°C	2 d 7 d	N/A
Sulphate	P/G	100 ml	Cool to $\leq 6^{\circ}$ C	28 d	АРНА
•	P/G			2 d	AS / ISO
Sulphite		100 ml	1ml EDTA/100ml (2.5g EDTA to 100ml)	2 d 7 d	
Sulphide (total)	P/G	100 ml	Cool + add 4 drops 2N Zinc Acetate/NaOH/100ml		APHA / ISO
Sulphide (diss)	P	100 ml	Filter & add 10ml Cu 2,4 DMP	12 hrs	AS
Sulphide (low level)	P	100 ml	pH 8-9.5 (NaOH) + Cool to $\leq 6^{\circ}$ C (potable water only)	7 d	AS
Tannins and Lignins	P/G	100 ml	Cool to <u><</u> 6°C	7 d	AS
Thiocyanate (SCN-)	P	100 ml	pH <2 (HNO ₃)	28 d	APHA
Thiosulphate	P/G	100 ml	Cool to <u><</u> 6°C	28 d	As per other Anions
TSS or TDS	P/G	200 ml	Cool to <u><</u> 6°C	7 d / 1 d	APHA / ISO
Turbidity	P/G	50 ml	Store in Dark	2 d	APHA
TVSS	Р	200ml	Cool to <u><</u> 6°C	1 d / 7d	USEPA / APHA
ORGANICS - V	OLATILE				
BTEX+ TRH C6-C9 (or TRH C6-C10)	Vial	2 x 40ml	pH <2 (H ₂ SO ₄ or HCl) or Sodium Bisulphate (NaHSO ₄)** + Cool to \leq 6°C	14 d	USEPA / AS
Haloacetic Acids	Vial	2 x 40ml	Sodium Bisulphate (NaHSO₄) + Cool to <u><</u> 6°C	14 d	USEPA
Trihalomethanes (THMs)	Vial	2 x 40ml	pH <2 (H ₂ SO ₄ or HCl) or Sodium Bisulphate (NaHSO ₄) + Cool to $\leq 6^{\circ}$ C	14 d	USEPA
VOCs (Brominated)	Vial	2 x 40 ml	− pH <2 (H₂SO₄ or HCl) or Sodium Bisulphate (NaHSO₄)** + Cool to <6°C	1 d	USEPA
VOCs (incl C1-C4 +	Vial	2 x 40ml	pH <2 (H_2SO_4 or HCI) or Sodium Bisulphate (NaHSO ₄)** +	7 d / 14 d	ISO / USEPA / AS
Methane, 1,4 Dioxane, Alcohols			Cool to <u><</u> 6°C		
Showing, Alcohols					
and Glycols) ORGANICS – S	SEMI VOLAT	<u> </u>			
and Glycols) ORGANICS – S	SEMI VOLAT	TILE 2 x 250 ml	Cool to <u>≺</u> 6°C (20mg sodium thiosulphate if residual chlorine)	30 d *	USEPA
and Glycols) ORGANICS — S Dioxins/Furans			Cool to <u><</u> 6°C (20mg sodium thiosulphate if residual chlorine) Cool to <u><</u> 6°C	30 d * 7 d	USEPA USEPA
and Glycols) ORGANICS – S Dioxins/Furans Diuron	G	2 x 250 ml			
and Glycols) ORGANICS – S Dioxins/Furans Diuron Explosives/Volatile	G G	2 x 250 ml 500 ml 200 ml (or 2 x 40 ml	Cool to <u><</u> 6°C	7 d	USEPA
and Glycols) ORGANICS – S Dioxins/Furans Diuron Explosives/Volatile Fatty Acids Glyphosate	G G	2 x 250 ml 500 ml 200 ml (or	Cool to <u><</u> 6°C	7 d	USEPA
and Glycols) DRGANICS – S Dioxins/Furans Diuron Explosives/Volatile Fatty Acids Slyphosate amily//Acrylamide	G G P/G	2 x 250 ml 500 ml 200 ml (or 2 x 40 ml VFAS) 100 ml	Cool to <u>≤</u> 6°C Cool to <u>≤</u> 6°C Cool to <u>≤</u> 6°C + Sodium Thiosulphate	7 d 7 d 14 d // 7 d	USEPA USEPA APHA / USEPA
and Glycols) ORGANICS – S Dioxins/Furans Diuron Explosives/Volatile Fatty Acids Glyphosate family//Acrylamide Diquat/Paraquat	G G G P/G P	2 x 250 ml 500 ml 200 ml (or 2 x 40 ml VFAS) 100 ml	Cool to $\leq 6^{\circ}$ C Cool to $\leq 6^{\circ}$ C Cool to $\leq 6^{\circ}$ C + Sodium Thiosulphate Cool to $\leq 6^{\circ}$ C	7 d 7 d 14 d // 7 d 7 d	USEPA USEPA APHA / USEPA USEPA
and Glycols) DRGANICS – S Dioxins/Furans Diuron Explosives/Volatile Fatty Acids Glyphosate amily//Acrylamide Diquat/Paraquat Ilicit Drugs	G G G P/G P G	2 x 250 ml 500 ml 200 ml (or 2 x 40 ml VFAs) 100 ml 500 ml 100mL	Cool to $\leq 6^{\circ}$ C	7 d 7 d 14 d // 7 d 7 d 7 d (estimate)	USEPA USEPA APHA / USEPA USEPA In house recommendation
and Glycols) DRGANICS – S Dioxins/Furans Diuron Explosives/Volatile Fatty Acids Glyphosate amily//Acrylamide Diquat/Paraquat Ilicit Drugs Pesticides or PCBs *	G G G P/G P G G	2 x 250 ml 500 ml 200 ml (or 2 x 40 ml VFAs) 100 ml 500 ml 100mL 2x100 ml^	Cool to $\leq 6^{\circ}$ C Cool to $\leq 6^{\circ}$ C Cool to $\leq 6^{\circ}$ C + Sodium Thiosulphate Cool to $\leq 6^{\circ}$ C Cool to $\leq 6^{\circ}$ C Cool to $\leq 6^{\circ}$ C	7 d 7 d 14 d // 7 d 7 d 7 d (estimate) 7 d *	USEPA USEPA APHA / USEPA USEPA In house recommendation USEPA
and Glycols) ORGANICS – S Dioxins/Furans Diuron Explosives/Volatile Fatty Acids Glyphosate family//Acrylamide Diquat/Paraquat Ilicit Drugs Pesticides or PCBs * Organometallics	G G G P/G P G G G	2 x 250 ml 500 ml 200 ml (or 2 x 40 ml VFAs) 100 ml 500 ml 2x100 ml ⁴ 500 ml	Cool to ≤6°C Cool to ≤6°C,	7 d 7 d 14 d // 7 d 7 d 7 d (estimate) 7 d * 7 d	USEPA USEPA APHA / USEPA USEPA In house recommendation USEPA ISO/USEPA
and Glycols) ORGANICS – S Dioxins/Furans Diuron Explosives/Volatile Fatty Acids Glyphosate family//Acrylamide Diquat/Paraquat Ilicit Drugs Pesticides or PCBs * Organometallics PBDEs	G G G P/G P G G G G	2 x 250 ml 500 ml 200 ml (or 2 x 40 ml VFAS) 100 ml 500 ml 2x100 ml ^A 500 ml 2x250 ml	Cool to $\leq 6^{\circ}$ C Cool to $\leq 6^{\circ}$ C, dark Cool to $\leq 6^{\circ}$ C, dark	7 d 7 d 14 d // 7 d 7 d 7 d (estimate) 7 d * 7 d 2 1 year	USEPA USEPA APHA / USEPA USEPA In house recommendation USEPA ISO/USEPA USEPA
and Glycols) ORGANICS – S Dioxins/Furans Diuron Explosives/Volatile Fatty Acids Glyphosate amily//Acrylamide Diquat/Paraquat Ilicit Drugs Pesticides or PCBs * Organometallics PBDEs Per- and	G G G P/G P G G G G G F	2 x 250 ml 500 ml 200 ml (or 2 x 40 ml VFAS) 100 ml 500 ml 2x100 ml ^A 500 ml 2x250 ml 2x250 ml	Cool to $\leq 6^{\circ}$ C Cool to $\leq 6^{\circ}$ C + Sodium Thiosulphate Cool to $\leq 6^{\circ}$ C Cool to $\leq 6^{\circ}$ C Cool to $\leq 6^{\circ}$ C Cool to $\leq 6^{\circ}$ C Cool to $\leq 6^{\circ}$ C, dark Cool to $\leq 6^{\circ}$ C, dark Cool to $\leq 6^{\circ}$ C, dark Cool to $\leq 6^{\circ}$ C	7 d 7 d 14 d // 7 d 7 d 7 d (estimate) 7 d * 7 d	USEPA USEPA APHA / USEPA USEPA In house recommendation USEPA ISO/USEPA
and Glycols) DRGANICS – S Dioxins/Furans Diuron Explosives/Volatile Fatty Acids Slyphosate amily//Acrylamide Diquat/Paraquat Ilicit Drugs Pesticides or PCBs * Drganometallics PBDEs Per- and Polyfluoroalkyl	G G G P/G P G G G G G P No Teflon liner	2 x 250 ml 500 ml 200 ml (or 2 x 40 ml VFAS) 100 ml 500 ml 100mL 2x100 ml ^A 500 ml 2x250 ml 2x60mL (std/low	Cool to $\leq 6^{\circ}$ C Cool to $\leq 6^{\circ}$ C Cool to $\leq 6^{\circ}$ C + Sodium Thiosulphate Cool to $\leq 6^{\circ}$ C Cool to $\leq 6^{\circ}$ C Cool to $\leq 6^{\circ}$ C Cool to $\leq 6^{\circ}$ C, dark Cool to $\leq 6^{\circ}$ C 2x250mL	7 d 7 d 7 d 14 d // 7 d 7 d 7 d (estimate) 7 d * 7 d ≥1 year 14 d / 28 d Longer holding times	USEPA USEPA APHA / USEPA USEPA In house recommendation USEPA ISO/USEPA USEPA
and Glycols) DRGANICS – S Dioxins/Furans Diuron Explosives/Volatile Fatty Acids Slyphosate amily//Acrylamide Diquat/Paraquat Ilicit Drugs Pesticides or PCBs * Drganometallics PBDEs Per- and Polyfluoroalkyl	G G G P/G P G G G G G F	2 x 250 ml 500 ml 200 ml (or 2 x 40 ml VFAS) 100 ml 500 ml 2x100 ml ^A 500 ml 2x250 ml 2x250 ml	Cool to $\leq 6^{\circ}$ C Cool to $\leq 6^{\circ}$ C + Sodium Thiosulphate Cool to $\leq 6^{\circ}$ C Cool to $\leq 6^{\circ}$ C Cool to $\leq 6^{\circ}$ C Cool to $\leq 6^{\circ}$ C Cool to $\leq 6^{\circ}$ C, dark Cool to $\leq 6^{\circ}$ C, dark Cool to $\leq 6^{\circ}$ C, dark Cool to $\leq 6^{\circ}$ C	7 d 7 d 14 d // 7 d 7 d (estimate) 7 d * 7 d ≥1 year 14 d / 28 d	USEPA USEPA APHA / USEPA USEPA In house recommendation USEPA ISO/USEPA USEPA
and Glycols) ORGANICS – S Dioxins/Furans Diuron Explosives/Volatile Fatty Acids Glyphosate family//Acrylamide Diquat/Paraquat Ilicit Drugs Pesticides or PCBs * Organometallics PBDEs Per- and Polyfluoroalkyl Substances (PFAS))	G G G P/G P G G G G G P No Teflon liner	2 x 250 ml 500 ml 200 ml (or 2 x 40 ml VFAS) 100 ml 500 ml 2x100 ml ^A 500 ml 2x250 ml 2x60mL (std/low leve/	Cool to $\leq 6^{\circ}$ C Cool to $\leq 6^{\circ}$ C Cool to $\leq 6^{\circ}$ C + Sodium Thiosulphate Cool to $\leq 6^{\circ}$ C Cool to $\leq 6^{\circ}$ C Cool to $\leq 6^{\circ}$ C Cool to $\leq 6^{\circ}$ C, dark Cool to $\leq 6^{\circ}$ C 2x250mL	7 d 7 d 7 d 14 d // 7 d 7 d 7 d (estimate) 7 d * 7 d ≥1 year 14 d / 28 d Longer holding times	USEPA USEPA APHA / USEPA USEPA In house recommendation USEPA ISO/USEPA USEPA / ASTM
and Glycols) ORGANICS – S Dioxins/Furans Diuron Explosives/Volatile Fatty Acids Glyphosate family//Acrylamide Diquat/Paraquat Illicit Drugs Pesticides or PCBs * Organometallics PBDEs Per- and Polyfluoroalkyl Substances (PFAS))	G G G P/G P G G G G G F No Teflon liner to be used	2 x 250 ml 500 ml 200 ml (or 2 x 40 ml VFAs) 100 ml 500 ml 2x100 ml ⁴ 500 ml 2x250 ml 2x250 ml 2x60mL (std/low level PFAS)	Cool to $\leq 6^{\circ}$ C Cool to $\leq 6^{\circ}$ C, dark Cool to $\leq 6^{\circ}$ C	7 d 7 d 7 d 14 d // 7 d 7 d 7 d (estimate) 7 d * 7 d ≥1 year 14 d / 28 d Longer holding times have been reported	USEPA USEPA APHA / USEPA USEPA In house recommendation USEPA ISO/USEPA USEPA / ASTM
and Glycols) ORGANICS – S Dioxins/Furans Diuron Explosives/Volatile Fatty Acids Glyphosate family//Acrylamide Diquat/Paraquat Illicit Drugs Pesticides or PCBs * Organometallics PBDEs Per- and Polyfluoroalkyl Substances (PFAS)) Pharmaceuticals/ Hormones Phenols – Speciated	G G G P/G P G G G G S S P No Teflon liner to be used G	2 x 250 ml 500 ml 200 ml (or 2 x 40 ml VFAS) 100 ml 500 ml 2x100 ml ^A 500 ml 2x250 ml 2x60mL (std/low level PFAS) 2x100mL	Cool to $\leq 6^{\circ}$ CCool to $\leq 6^{\circ}$ C, darkCool to $\leq 6^{\circ}$ C, darkCool to $\leq 6^{\circ}$ C2x250mL(trace level PFAS)Cool to $\leq 6^{\circ}$ C	7 d 7 d 7 d 14 d // 7 d 7 d 7 d (estimate) 7 d * 7 d ≥1 year 14 d / 28 d Longer holding times have been reported 7d (estimate)	USEPA USEPA APHA / USEPA USEPA In house recommendation USEPA USEPA USEPA / ASTM
and Glycols) ORGANICS – S Dioxins/Furans Diuron Explosives/Volatile Fatty Acids Glyphosate amily//Acrylamide Diquat/Paraquat Ilicit Drugs Pesticides or PCBs * Organometallics PBDEs Per- and Polyfluoroalkyl Substances (PFAS)) Pharmaceuticals/ Hormones Phenols – Speciated Phenoxy Herbs	G G G P/G P G G G G G No Teflon liner to be used G G	2 x 250 ml 500 ml 200 ml (or 2 x 40 ml VFAS) 100 ml 500 ml 100mL 2x100 ml ^A 500 ml 2x250 ml 2x250 ml (std/low level PFAS) 2x100mL 2x100 ml ^A	Cool to $\leq 6^{\circ}$ CCool to $\leq 6^{\circ}$ C + Sodium ThiosulphateCool to $\leq 6^{\circ}$ CCool to $\leq 6^{\circ}$ CCool to $\leq 6^{\circ}$ CCool to $\leq 6^{\circ}$ CCool to $\leq 6^{\circ}$ C, darkCool to $\leq 6^{\circ}$ C, darkCool to $\leq 6^{\circ}$ C2x250mL(trace level PFAS)Cool to $\leq 6^{\circ}$ CCool to $\leq 6^{\circ}$ C	7 d 7 d 7 d 14 d // 7 d 7 d 7 d (estimate) 7 d * 7 d ≥1 year 14 d / 28 d Longer holding times have been reported 7 d (estimate) 7 d *	USEPA USEPA APHA / USEPA USEPA USEPA USEPA USEPA / ASTM USEPA / ASTM
and Glycols) ORGANICS – S Dioxins/Furans Diuron Explosives/Volatile Fatty Acids Glyphosate amily//Acrylamide Diquat/Paraquat Ilicit Drugs Pesticides or PCBs * Organometallics PBDEs Per- and Polyfluoroalkyl Substances (PFAS)) Pharmaceuticals/ Hormones Phenols – Speciated Phenoxy Herbs	G G G P/G P G G G G No Teflon liner to be used G G G	2 x 250 ml 500 ml (or 2 x 40 ml VFAS) 100 ml 500 ml 100mL 2x100 ml ^A 500 ml 2x250 ml 2x60mL (<i>std/low</i> <i>level</i> <i>PFAS</i>) 2x100 ml ^A 2x100 ml ^A	Cool to $\leq 6^{\circ}$ CCool to $\leq 6^{\circ}$ C, darkCool to $\leq 6^{\circ}$ C, darkCool to $\leq 6^{\circ}$ C, darkCool to $\leq 6^{\circ}$ C2x250mL(trace level PFAS)Cool to $\leq 6^{\circ}$ CCool to $\leq 6^{\circ}$ CCool to $\leq 6^{\circ}$ CCool to $\leq 6^{\circ}$ CCool to $\leq 6^{\circ}$ C	7 d 7 d 7 d 14 d // 7 d 7 d 7 d (estimate) 7 d * 7 d ≥1 year 14 d / 28 d Longer holding times have been reported 7 d (estimate) 7 d * 7 d *	USEPA USEPA APHA / USEPA USEPA IN house recommendation USEPA USEPA USEPA / ASTM IN house recommendation USEPA USEPA
and Glycols) ORGANICS – S Dioxins/Furans Diuron Explosives/Volatile Fatty Acids Glyphosate family//Acrylamide Diquat/Paraquat Illicit Drugs Pesticides or PCBs * Organometallics PBDEs Per- and Polyfluoroalkyl Substances (PFAS)) Pharmaceuticals/ Hormones Phenols – Speciated Phenoxy Herbs Phenoxy Herbs Phenoxy Herbs	G G G P/G P G G G G P No Teflon liner to be used	2 x 250 ml 500 ml (or 2 x 40 ml VFAS) 100 ml 500 ml 100mL 2x100 ml ^A 500 ml 2x250 ml 2x60mL (<i>std/low</i> <i>level</i> <i>PFAS</i>) 2x100 ml ^A 2x250 ml ^A 2x250 ml ^A	Cool to $\leq 6^{\circ}$ CCool to $\leq 6^{\circ}$ C, darkCool to $\leq 6^{\circ}$ C, darkCool to $\leq 6^{\circ}$ C, darkCool to $\leq 6^{\circ}$ C2x250mL(trace level PFAS)Cool to $\leq 6^{\circ}$ CCool to $\leq 6^{\circ}$ C	7 d 7 d 7 d 14 d // 7 d 7 d (estimate) 7 d * 7 d ≥1 year 14 d / 28 d Longer holding times have been reported 7 d (estimate) 7 d * 7 d * 14 d / 28 d	USEPA USEPA APHA / USEPA USEPA USEPA USEPA USEPA / ASTM In house recommendation USEPA / ASTM
and Glycols)	G G G P/G P G G G G V No Teflon liner to be used G G G G G G	2 x 250 ml 500 ml 200 ml (or 2 x 40 ml VFAs) 100 ml 500 ml 2x100 ml ^A 500 ml 2x250 ml 2x60mL (std/low level PFAS) 2x100 ml ^A 2x100 ml ^A 2x250 ml 2x100 ml ^A	Cool to $\leq 6^{\circ}$ CCool to $\leq 6^{\circ}$ C, darkCool to $\leq 6^{\circ}$ C2x250mL(trace level PFAS)Cool to $\leq 6^{\circ}$ CCool to $\leq 6^{\circ}$ CCool to $\leq 6^{\circ}$ CCool to $\leq 6^{\circ}$ CpH <2 (H ₂ SO ₄ or HCl) + Cool to $\leq 6^{\circ}$ C	7 d 7 d 7 d 14 d // 7 d 7 d (estimate) 7 d * 7 d ≥1 year 14 d / 28 d Longer holding times have been reported 7 d (estimate) 7 d * 7 d * 7 d * 7 d 28 d	USEPA USEPA APHA / USEPA USEPA USEPA ISO/USEPA USEPA / ASTM USEPA / ASTM USEPA USEPA USEPA USEPA USEPA USEPA
and Glycols) ORGANICS – S Dioxins/Furans Diuron Explosives/Volatile Fatty Acids Glyphosate family//Acrylamide Diquat/Paraquat Illicit Drugs Pesticides or PCBs * Organometallics PBDEs Per- and Polyfluoroalkyl Substances (PFAS)) Pharmaceuticals/ Hormones Phenols – Speciated Phenoxy Herbs PAHs or Phthalates SVOCs including – OCPs, OPPs, PCBs,	G G G P/G P G G G G V No Teflon liner to be used G G G G G G	2 x 250 ml 500 ml 200 ml (or 2 x 40 ml VFAS) 100 ml 500 ml 100mL 2x100 ml ^A 500 ml 2x250 ml 2x250 ml (std/low level PFAS) 2x100mL 2x100 ml ^A 2x250 ml 2x100 ml ^A 2x250 ml 2x100 ml ^A 2x100 ml ^A 2x100 ml ^A	Cool to $\leq 6^{\circ}$ CCool to $\leq 6^{\circ}$ C, darkCool to $\leq 6^{\circ}$ C, darkCool to $\leq 6^{\circ}$ C, darkCool to $\leq 6^{\circ}$ C2x250mL(trace level PFAS)Cool to $\leq 6^{\circ}$ CCool to $\leq 6^{\circ}$ C	7 d 7 d 7 d 14 d // 7 d 7 d (estimate) 7 d * 7 d ≥1 year 14 d / 28 d Longer holding times have been reported 7 d (estimate) 7 d * 7 d * 7 d * 7 d 28 d	USEPA USEPA APHA / USEPA USEPA USEPA ISO/USEPA USEPA / ASTM USEPA / ASTM USEPA USEPA USEPA USEPA USEPA USEPA
and Glycols) ORGANICS – S Dioxins/Furans Diuron Explosives/Volatile Fatty Acids Glyphosate family//Acrylamide Diquat/Paraquat Illicit Drugs Pesticides or PCBs * Organometallics PBDEs Per- and Polyfluoroalkyl Substances (PFAS)) Pharmaceuticals/ Hormones Phenols – Speciated Phenoxy Herbs Phenoxy Herbs PAHs or Phthalates SVOCs including –	G G G P/G P G G G G V No Teflon liner to be used G G G G G G	2 x 250 ml 500 ml 200 ml (or 2 x 40 ml VFAS) 100 ml 500 ml 100mL 2x100 ml ^A 500 ml 2x250 ml 2x50 ml (std/low level PFAS) 2x100 ml ^A 2x100 ml ^A 2x250 ml 2x100 ml ^A	Cool to $\leq 6^{\circ}$ CCool to $\leq 6^{\circ}$ C, darkCool to $\leq 6^{\circ}$ C, darkCool to $\leq 6^{\circ}$ C, darkCool to $\leq 6^{\circ}$ C2x250mL(trace level PFAS)Cool to $\leq 6^{\circ}$ CCool to $\leq 6^{\circ}$ C	7 d 7 d 7 d 14 d // 7 d 7 d (estimate) 7 d * 7 d ≥1 year 14 d / 28 d Longer holding times have been reported 7 d (estimate) 7 d * 7 d * 7 d * 7 d 28 d	USEPA USEPA APHA / USEPA USEPA USEPA USEPA USEPA / ASTM USEPA / ASTM USEPA / ASTM USEPA USEPA USEPA USEPA USEPA

Page 2 of 8



Test	Bottle	Volume	Preservation (Bottles are colour coded, see labels)	RHT	Reference
MICROBIOLO	GICAL and	ALGAE			
Acid Producing or	P/G (Sterile	100 ml	Cool to <6°C + Sodium Thiosulphate	1 d	AS
Denitrifying Bacteria					
Amoeba	P (Sterile)	250mL	Room Temperature e.g. 20ºC±5ºC ideally	3d	AS
Legionella	P (Sterile)	100mL	Cool to <u><</u> 6°C + Sodium Thiosulphate	2d	AS
Heterotrophic Plate	P (Sterile)	100mL	Cool to <u><</u> 6°C + Sodium Thiosulphate	1d	APHA
Count					
TT.coli/E.coli, T.coli,	P (Sterile)	100mL	Cool to <u><</u> 6°C + Sodium Thiosulphate	1d	AS
Faecal.Enterococci,		(each)			
Pseudo, Clostridium					
Chlorophyll-a, b, c &	P/G	500 ml	Cool <u><</u> 6°C + Dark or Filter/ Freeze	2 d / 28 d	АРНА
Phaeophytin					
Micro/Algae in	P (Sterile)	250-500ml	Cool to <u>≺</u> 6°C	1 d	AS
natural water					
Micro/Algae in tap	P (Sterile)	250-500ml	Cool to <u><</u> 6°C + Sodium Thiosulphate	1 d	AS
water					
Iron Related	P (Sterile)	100ml	Sterile, Sodium Thiosulphate + Cool to \leq 6°C . No headspace	1 d	AS
Bacteria					
Sulphate Reducing	P (Sterile)	100ml	Sterile, NO Sodium Thiosulphate + Cool to \leq 6°C . No	1 d	AS
Bacteria			headspace		
RADIOLOGIC	<u>AL</u>				
Radiochem – Gross	Р	200 mL	pH <2 HNO₃	30 d	ISO
Alpha/Beta					
Radiochem – Gross	Р	1000 ml	none	7 d	ISO
Alpha/Beta					
Radiochem –	Р	1000 ml	pH <2 (HNO₃)	1 m	ISO
Radium 226 + 228					

Water Comments:

P/G: Plastic/Glass respectively

* Dioxins - 30 days till extraction. 45 days after extraction. * Semi-volatiles - 7 days till extraction, 40 days after extraction.

* THM's – add 10% thiosulfate if residual chlorine present. * TRH (C10-C36 or C10-C40) – 7 days if treated as semi-volatile.

** Use Sodium Bisulphate where Dangerous Goods restrictions apply.

^ An absolute minimum volume of 100mL for SVOC analysis is required. Where low/trace levels are required 2 x 250mL is recommended.

^{\$} Chlorite is sub-contracted.

[®] Perchlorate can be unpreserved if Oxyhalides are not required

VOC/vTRH/THM - A third vial is preferred for VOC waters to allow for confirmation (if required) and QC sample analysis (duplicates and spikes).

SVOC/svTRH – A third bottle (100mL or 250mL for low level analysis) is preferred for SVOC waters to allow for confirmation (if required) and QC sample analysis (duplicates and spikes).

For specialised and/or boutique Organics, please contact the laboratory directly if recommended THTs are not clear from the information above.

For dissolved metals (such as ground waters) samples should be 0.45µm filtered into a HNO₃ preserved bottle. For dissolved metals that are not field filtered and acidified, Envirolab will filter on receipt from an unpreserved sample container then acidify. There will be a filtration charge and some elements may be affected by this process. For total recoverable metals (such as surface waters) samples should be placed unfiltered into a HNO₃ preserved bottle. For total recoverable metals that are not field preserved, Envirolab will sub sample from the unpreserved bottle on receipt and acidify with HNO₃ prior to sample digestion.

For Ferrous Iron (Fe²⁺) samples should be 0.45µm filtered into a HCl preserved bottle. Discrepancies may occur if not field filtered.

For Hexavalent Chromium (Cr⁶⁺) samples should be 0.45µm filtered into a NaOH preserved bottle. Discrepancies may occur if not field filtered.

The above preservation techniques have been chosen by the Envirolab Group as suitable for most situations. There are other preservation techniques available – please contact Envirolab Customer Service for other options.

References:

AS	=	Australian Standard 5667.1:1998 Water Quality Sampling or latest version
APHA	=	Standard Methods for the Examination of Water & Wastewater, Latest Edition
ISO	=	ISO 5667.3 2012 or latest version
USEPA	=	USEPA SW846 or latest version
NEPM	=	Schedule B(3) 2013 or latest version



RECOMMENDED PRESERVATION & HOLDING TIMES (RHT) FOR SOILS AND SEDIMENTS

www.envirolab.com.au

ADELAIDE: (08) 7087 6800 BRISBANE: (07) 3266 9532

DARWIN: 0477 012 027 PERTH: (08) 9317 2505

MELBOURNE (03) 9763 2500	SYDNE	Y: (02) 9910 6200	PERTH: (08) 9317 2505
Test	Jar	Quantity	Preservation	RHT	Reference
INORGANICS, ME	TALS AND PHYSIC	AL TESTS			
AMD	Bag	1-2 kg	Nil	N/A	-
Asbestos	Bag	50 g	Nil	N/A	AS4964
Asbestos (WA guidelines)	Bag	500 mL jar	Nil	N/A	NEPM
ASS /AVS	Bag	100 g	Remove air and freeze (Ziplock Bag)	<24 hr#	QASSMAC
Bromide / Iodide**	P/G	20 g	Cool to <u><</u> 6°C	28 d	NEPM
Cation Exchange Capacity	G	50 g	Cool to <u><</u> 6°C	28 d	NEPM
Carbon – TOC	G	20 g	Cool to <u><</u> 6°C, no headspace	28 d	NEPM
Chloride	P/G	20 g	Cool to <u><</u> 6°C	28 d	NEPM
Clay	P/G/Bag	200 g	Nil	Not Determined	
Conductivity	P/G	20 g	Cool to <u><</u> 6°C	7 d	NEPM
Cyanates	P/G	20 g	Cool to <u><</u> 6°C, dark	1 d	In House
Cyanides	P/G	20 g	Cool to <u><</u> 6°C, dark	14 d	NEPM
Explosives	G	20 g	Cool to <u><</u> 6°C	14 d	USEPA
Field Peroxide	Bag	100 g	Remove air and freeze (Ziplock Bag)	<24 hr	QASSMAC
Foreign Material	Bag	6 kg	Nil	N/A	N/A
Fluoride	P/G	20 g	Cool to <u><</u> 6°C	28 d	NEPM
Hexavalent Cr	P/G	20 g	Cool to $\leq 6^{\circ}$ C	28 d	USEPA
LEAF 1313 (variable pH)	P/G	1-2kg	Cool to <u><</u> 6°C	As per analyte of interest	USEPA
LEAF 1314 (column test)	P/G	2-3kg	Cool to <u><</u> 6°C	As per analyte of interest	USEPA
LEAF 1315 (monolith test)	P/G	2-3kg	Cool to <u><</u> 6°C	As per analyte of interest	USEPA
LEAF 1316 (variable L/S ratio) P/G	2-5kg	Cool to <u><</u> 6°C	As per analyte of interest	USEPA
Metals (except CrVI, Hg)	P/G	20 g	Cool to <u><</u> 6°C	6 m	USEPA
Mercury	P/G	20 g	Cool to <u><</u> 6°C	28 d	USEPA
Moisture	P/G	20 g	Cool to <u><</u> 6°C	14 d	NEPM
Nutrients – Ammonia, TKN, 1	rp G	20 g	Cool to <u><</u> 6°C	28 d	USEPA
Nutrients – TN, Nitrate, Nitri		20 g		7 d	In house
Phosphate			Cool to <u><</u> 6°C		recommendation
рH	P/G	20 g	Cool to <u><</u> 6°C	7 d	NEPM
Particle size Distribution (PSI	D) Bag	1 kg (sand/clay), 2 kg (gravel)	Nil	None	-
Sulphate	P/G	20 g	Cool to <u><</u> 6°C	28 d	NEPM
SPOCAS	P	100 g	Remove air and freeze (Ziplock Bag)	<24 hr#	QASSMAC
Sulphide	P/G	20 g	Cool to <6°C + 2M Zn acetate to cover	7 d	NEPM
TCLP	P/G	100 g	 Cool to <u><</u> 6°C	As per analyte of interest	
Thiocyanates	P/G	20 g	 Cool to <u><</u> 6°C, dark	14 d	In House
ORGANICS – SEMI		Ŭ			
Dioxins and Furans	G	250 g	Cool to <u><</u> 6°C, dark	30 d	NEPM
Herbicides	G	20 g	Cool to <u><</u> 6°C	14 d	USEPA
Illicit Drugs	G	50g	Cool to <u><</u> 6°C	28 d (estimate)	In house
Organotins	G	20 g	Cool to <u><</u> 6°C	14 d	NODG
OC/OP's	G	20 g	Cool to <u><</u> 6°C	14 d	USEPA
PAH's	G	20 g	Cool to <u><</u> 6°C	14 d	USEPA
PCB's	G	20 g	Cool to <u><</u> 6°C, dark	28 d	USEPA
Per- and Polyfluoroalkyl Substances (PFAS inc PFOS/PFOA/PFHxS)	Р	20 g	Cool to <u><</u> 6°C	60 d / 28 d Longer holding times have been reported	USEPA / ASTM
Phenols – Speciated (inc Coa	l Tar) G	20 g	Cool to <u><</u> 6°C	14 d	USEPA
SVOC's	G	20 g	Cool to <u><</u> 6°C	14 d	USEPA

TRH (C_{10⁻40}) and Speciated

G

20 g

Cool to <u><</u>6°C

USEPA

14 d



Test	Jar	Qty	Preservation	RHT	Reference
SOILS (cont.)					
ORGANICS - VOLATILE					
BTEX+C ₆ -C ₁₀	G	20 g	Cool to <u><</u> 6°C, no headspace	14 d	USEPA
zHE TCLP	P/G	100 g		As per analytes of	
				interest	
TRH (C ₆₋₉ or C ₆₋₁₀)	G	20 g	Cool to <6°C, no headspace	14 d	USEPA
VOCs / VHCs / THMs / VACs	G	20 g	Cool to <6°C, no headspace	14 d *	USEPA / NEPM
/1, 4 Dioxane / Alcohols / Glycols					
MICROBIOLOGICAL					
Micro (E.Coli, FC, TC etc.)	P (sterile)	125 g	Cool to <u><</u> 6°C	<24 hr	USEPA
SEDIMENTS					
Elutriate Testing	P/G	100-500 g (depending on		14 d	NAGD
		analytes) + supply of 5L of	Cool to <u><</u> 6°C, dark, airtight		
		seawater			
Metals (except for Hg)	P/G	20 g	Cool to <6°C / freeze for extended storage	6 months	NAGD/NODG
Metals – Hg only			Cool to <6°C/ freeze for extended storage	28 d	NAGD/NODG
Organics (e.g. PCB's, Pesticides,	G	20 g	Cool to <6°C, dark/ freeze for extended	14 d	NAGD/NODG
PAH and TBT)			storage		

P/G: Plastic/Glass respectively

250mL jar ≈ 350-420g

125mL jar ≈ 175-210g

Soil Comments:

- One 250ml size jar will cover most determinations in soil (extra is preferable for sizing tests e.g. Clay determination).
- Asbestos should be sampled in its own plastic bag and should be about 1 large handful or about 40g.
- #SPOCAS/CrS should be supplied in its own plastic bag, air removed and frozen if possible or delivered to the laboratory within 24hrs of sampling.
- * Less for vinyl chloride, styrene, 2-chloroethyl vinyl ether (7 days)
- Leachate Environmental Assessment Framework (*LEAF*) quantities can be project specific, please contact the laboratory for advice.
- ** Some analytes e.g. lodide do not have recommended holding times, hence holding times for similar analytes may have been recommended.

Sediment Comments:

- Where samples are to be subsampled or homogenised in the lab, they should be stored at 2-8°C in the dark and transported to the laboratory within 72 hours (preferably 24 hours) of collection (NODG 2002)
- For sediment samples containing TBT, mercury and other volatiles for chemical analysis, freezing to below -10°C within 12 hours of collection, and before their despatch in a well-insulated cooler, improves laboratory storage life (NODG 2002).

References:

- USEPA = USEPA SW846
- NEPM = Schedule B(3) 2013
- NODG = National Ocean Disposal Guidelines 2002
- NAGD = National Assessment Guidelines for Dredging 2009

Soil samples for agricultural tests can be collected in zip lock plastic bags or glass jars, the literature suggests that there is evidence that field moist soils can be kept at $\leq 6^{\circ}$ C for days or weeks before any change in nutrient cations and anions. Air-drying and/or freezing will assist in reducing the potential for biological transformation. Hence analysis should typically occur within two weeks of sampling.



RECOMMENDED PRESERVATION & HOLDING TIMES (RHT) FOR AIR

www.envirolab.com.au

ADELAIDE:	(08) 7087 6800	BRISBANE: (07) 3266 9532	DARWIN: 0477 012 027
MELBOURNE	(03) 9763 2500	SYDNEY: (02) 9910 6200	PERTH: (08) 9317 2505

Determination	Container	Preservation	RHT	Reference
Acid Gases/Mists (HCl, HNO3, H2SO4, H3PO4, HBr)	Sorbent / Filter	Cool, <u><</u> 6°C	21 d sorbents / 28 d filters	NIOSH 7903 (sorbents) / NIOSH 7907-7908 (filters)
Ammonia	Sorbent / Filter	Cool, <u><</u> 6°C	21 d sorbents / 14 d filters	NIOSH / best practice
Diesel Particulates	Filter	Ambient	180 d	NIOSH 5040
Fluoride	Filter / Impinger	Cool, <u><</u> 6°C	28 d/best practice	NIOSH 7906/USEPA m13b
Halogens (Br ₂ /Cl ₂)	Sorbent	Cool, <u><</u> 6°C	30 d	NIOSH 6011
NO/NO2	Sorbent	Cool, <u><</u> 6°C	7 d	NIOSH 6014
Hydrogen Cyanide	Sorbent	Cool, <u>≤</u> 6°C	14 d (sorbent) / ASAP best practice (passive)	NIOSH 6010/OSHA 1015
Hydrogen Sulphide (or see Sulphur Compounds below)	Sorbent	Cool, <u><</u> 6°C	30 d	NIOSH 6013
Sulphur Dioxide – SO ₂	Filter/Sorbent	Cool, <u><</u> 6°C	30 d (best practice)	NIOSH 6004
SOx by USEPA m6, 8	Impinger	Cool, <u><</u> 6°C	30 d	USEPA m6 & m8
Metals general (includes CrVI)	Filters / Sorbents	Cool, <u><</u> 6°C	180 d (14 d for CrVI)	NIOSH 7303 (Metals) / NIOSH 7600 (Cr6+)
Metals by USEPA m29	Filters / Impingers	Cool, <u><</u> 6°C	180 d (best practice)	USEPA m29
Formaldehyde	Sorbents	Cool, <u><</u> 6°C	35 d	NIOSH 2016
Organic Vapours (includes Radiello)	Sorbents	Cool, <u><</u> 6°C	30 d (best practice)	OSHA/NIOSH
Volatile Organics* by EPA TO-15	Canister	Ambient	30 d	TO-15
Volatile Organics* by EPA TO-15, modified	Air Sampling (Tedlar) bag	Ambient/Dark	72 hrs	TO-15
Volatile Organics* by EPA TO-17 (includes Radiello)	Sorbent tube	Cool, <u><</u> 6°C	30 d	TO-17
Air Petroleum Hydrocarbons by APH method	Canister	Ambient	30 d	Air-Phase Petroleum Hydrocarbons (APH), 2009 & TO-15
Air Petroleum Hydrocarbons by APH method (includes Radiello)	Sorbent tube	Ambient	30 d	Air-Phase Petroleum Hydrocarbons (APH), 2009 & TO-15
Air Petroleum Hydrocarbons by APH method	Air Sampling (Tedlar) bag	Ambient/Dark	72 hrs	Air-Phase Petroleum Hydrocarbons (APH), 2009 & TO-15
Gases by ASTM1945, ASTM1946 & EPA 3c modified as listed below (includes C1- C6)	Canister	Ambient	30 d	ASTM1945, ASTM1946 & EPA 3c
Gases by ASTM1945, ASTM1946 & EPA 3c modified as listed below (includes C1- C6)	Air Sampling (Tedlar) bag	Ambient	72 hrs	ASTM1945, ASTM1946 & EPA 3c
Reduced Sulphur Compounds *	Air Sampling (Tedlar) bag	Ambient	24 hrs	ASTM D5504
Reduced Sulphur Compounds *	Canister (Silico treated)	Ambient	72 hrs	ASTM D5504
Aldehydes and Ketones	Sorbent tube	Cool, <u><</u> 6°C	14 d	USEPA TO-11A



Determination	Container	Preservation	RHT	Reference
Polynuclear Aromatic Hydrocarbons	PUF/XAD	Cool, <u><</u> 6°C	7 d (or 14d [#])	USEPA TO-13A
Organochlorine Pesticides and PCBs	PUF	Cool, <u><</u> 6°C	7 d	USEPA TO-4A USEPA TO-10A
Dioxins/Furans	XAD	Cool, <u><</u> 6°C	30 d	In house recommendation

* Analyte integrity may be affected by high humidity, particularly for oxygenated and reduced sulphur compounds.

As air sampling bags have a short recommended holding period, it is advisable the laboratory is contacted prior to sampling to ensure analysis occurs within the recommended holding time.

Air Sampling (Tedlar) bags should be partially filled when shipped by air freight.

Gases by ASTM1945, ASTM1946 & EPA 3c includes CH_4 , CO, CO₂, He, H₂, N₂, O₂ and >20 C₂-C₆ Aliphatic Hydrocarbons

[#]Canadian government laboratory reported stability for up to 14 days for PAHs

Some references have become obsolete (e.g. some NIOSH methods) and are therefore indicative only.

For particular sorbents and/or samplers (e.g. SKC, Radiello, 3M and Waterloo Membrane Samplers), please see manufacturer specific information, generally available online.

RECOMMENDED PRESERVATION & HOLDING TIMES (RHT) FOR BIOLOGICAL SAMPLES

www.envirolab.com.au

ADELAIDE:	(08) 708	37 6800	BRISBAN	IE: (07) 3266 9532	DARWIN:	0477 012 027
MELBOURNE	(03) 976	3 2500	SYDNE	Y: (02) 9910 6200	PERTH:	(08) 9317 250
Test	Co	ontainer	Quantity	Preservation	RHT	Reference
BLOOD/PLASM	A/SERUM					
PFAS/TOPA	S	erum eparating	1 x 8mL SST gold cap	Cool to <u><</u> 6°C SST for Serum	2 months	USEPA (adopted draft soil
	Т	ube (SST)	[1 x 4mL purple cap with anti-coagulant]	[K₂EDTA (blood only – non-routine request)]		criteria)
URINE			5 -	·		
Metals						
Metals including M	ercury U	lrine jars	20mL specimen jar	Cool to <u><</u> 6°C	28	NIOSH
Speciated Arsenic	U	lrine jars	20mL specimen jar	Filter, Cool to <u><</u> 6°C /freeze	2 months	*Reference
						Paper
Organics						
BTEX Metabolites e S-PMA	.g. U	lrine jars	20mL specimen jar	Cool to <u><</u> 6°C/frozen, dark	7/30 days	NIOSH
Hexane Metabolite 2,5-HD	se.g. U	lrine jars	20mL specimen jar	Cool to \leq 6°C/frozen, dark	7/30 days	NIOSH
PAH Metabolites e.	g. 1-HP U	lrine jars	20mL specimen jar	Cool to <u><</u> 6°C/frozen, dark	7/30 days	NIOSH
Phenoxy Acid Herbi	cides U	lrine jars	20mL specimen jar	Cool to <u><</u> 6°C/frozen, dark	7/30 days	NIOSH
Speciated Phenols	U	lrine jars	20mL specimen jar	Cool to <u><</u> 6°C/frozen, dark	7/30 days	NIOSH
VOC Metabolites e. Mandelic acid	g. U	Irine jars	20mL specimen jar	Cool to <u><</u> 6°C/frozen, dark	7/30 days	NIOSH

References:

 * Sample Preparation and Storage Can Change Arsenic Speciation in Human Urine – Feldmann et al Clinical Chemistry 45, No. 11, 1999



RECOMMENDED PRESERVATION & HOLDING TIMES (RHT) FOR ILLICIT DRUGS on SWABs

Test	Container	Quantity	Preservation	RHT	Reference
ILLICIT DRUGS					
Methamphetamine and	Swab kit –	2 swabs per 100cm ²	Cool to <u><</u> 6°C	30 days	NIOSH
three precursors: MDMA,	50mL tube				
Ephedrine and					
Pseudoephedrine					
Other analyse available e.g.					
Cocaine, Heroin, Fentanyl,					
Cannabinoids.					